

THE TRANSFORMATION GROUPS WHOSE C^* -ALGEBRAS ARE FELL ALGEBRAS

ASTRID AN HUEF

ABSTRACT. Let (G, X) be a locally compact transformation group, in which G acts freely on X . We show that the associated transformation-group C^* -algebra $C_0(X) \rtimes G$ is a Fell algebra if and only if X is a Cartan G -space.

An irreducible representation π of a C^* -algebra A is a *Fell point* of the spectrum $\hat{A}$ if there exist a neighbourhood V of π in $\hat{A}$ and an element a of the positive cone A^+ of A such that $\sigma(a)$ is a rank-one projection for all $\sigma \in V$. If every irreducible representation is a Fell point then A is a *Fell algebra*. A C^* -algebra has *continuous trace* if and only if its spectrum is Hausdorff and it is a Fell algebra.

Let (G, X) be a locally compact Hausdorff transformation group: thus G is a locally compact Hausdorff group and X is a locally compact Hausdorff space together with a jointly continuous map $(s, x) \mapsto s \cdot x$ from $G \times X$ to X such that $s \cdot (t \cdot x) = st \cdot x$ and $e \cdot x = x$. We assume that both G and X are second countable. The action of G on X lifts to a strongly continuous action α of G by automorphisms of $C_0(X)$ given by $\alpha_s(f)(x) = f(s^{-1} \cdot x)$. The associated transformation-group C^* -algebra $C_0(X) \rtimes G$ is the C^* -algebra which is universal for the covariant representations of the C^* -dynamical system $(C_0(X), G, \alpha)$ in the sense of [10]. More concretely, $C_0(X) \rtimes G$ is the enveloping C^* -algebra of the Banach $*$ -algebra $L^1(G, C_0(X))$ of functions $f : G \rightarrow C_0(X)$ which are integrable with respect to a fixed left Haar measure μ on G [8, Section 7.6], with multiplication and involution given by

$$f * g(s) := \int_G f(r) \alpha_r(g(r^{-1}s)) \, d\mu(r) \quad \text{and} \quad f^*(s) := \Delta(s^{-1}) \alpha_s(f(s^{-1}))^*,$$

where Δ is the modular function associated with μ .

In [3] Green showed that if the action of G on X is free, then $C_0(X) \rtimes G$ has continuous trace if and only if the action is proper. The purpose of this note is to obtain a parallel characterisation of the transformation groups whose C^* -algebras are Fell algebras.

Theorem 1. *Let (G, X) be a second countable transformation group with G acting freely on X . Then $A = C_0(X) \rtimes G$ is a Fell algebra if and only if X is a Cartan G -space.*

A subset N of X is *wandering* if the set $\{s \in G : s \cdot N \cap N \neq \emptyset\}$ is relatively compact in G . If every point of X has a wandering neighbourhood then X is a *Cartan G -space* [7, Definition 1.1.2]. When the action is free, these spaces are precisely the principal bundles of Cartan by [7, Theorem 1.1.3], and they are called principal G -bundles by Husemoller in [5]. The action

Date: July 23, 1999.

This paper is based on part of the author's doctoral dissertation written at Dartmouth College. We thank Dana Williams for his supervision and help and Marc Rieffel for a suggestion regarding the exposition of this paper.

of G on X is *proper* (in the sense that the map $(s, x) \mapsto (s \cdot x, x)$ from $G \times X$ into $X \times X$ is a proper map) if and only if every compact subset of X is wandering. If the action of G on X is proper then X is a Cartan G -space, but the converse is not true even if the action is free (see Example 3).

For the proof of Theorem 1 we require some background material. In [1], Archbold and Somerset characterised Fell algebras A as follows: define S to be the set of all $a \in A^+$ such that

$$\mathrm{tr}(\pi_\alpha(a)) \rightarrow \sum_{\pi \in L} \mathrm{tr}(\pi(a)) < \infty \quad (\dagger)$$

whenever $\{\pi_\alpha\}$ is a properly convergent net in $\hat{A}$ with limit set L . (A properly convergent net is a convergent net which converges to every cluster point.) The linear span $\mathcal{S}(A)$ of S is a two-sided (hereditary) ideal of A , and A is a Fell algebra if and only if $\mathcal{S}(A)$ is dense in A [1, Corollary 3.7]. To prove Theorem 1 we show that if X is not a Cartan G -space then there is an element in the Pedersen ideal of an ideal I of $C_0(X) \rtimes G$ which is not an element of $\mathcal{S}(I)$. This suffices because an ideal of a Fell algebra is itself Fell, and because the Pedersen ideal is minimal among dense ideals [8, Theorem 5.5.1].

We thank Robert Archbold for pointing out that we can use [6, Theorem 3] instead of [2, Theorem 2.6 and Proposition 2.1 i)] in the proof below.

Proof of Theorem 1. We prove the easier direction first. If X is a Cartan G -space then the orbits are closed by [7, Proposition 1.1.4] and thus A is CCR [11]. Because the action is free the map $x \mapsto \pi_x := \mathrm{Ind}_{\{e\}}^G(\epsilon_x)$ induces a homeomorphism of X/G onto $\hat{A}$ [3, Lemma 16]. Fix $x \in X$ and let U be an open wandering neighbourhood of x in X . The action of G on $G \cdot U$ is proper by [7, Proposition 1.2.4]. Because the action is free the algebra $C_0(G \cdot U) \rtimes G$ has continuous trace [3, Theorem 17], and since $C_0(G \cdot U)$ is a G -invariant ideal of $C_0(X)$ this crossed product embeds naturally as an ideal J in $C_0(X) \rtimes G$ [3, Lemma 16]. Viewing π_x as a representation of J shows that it is a Fell point of the open subset $\hat{J}$ of $\hat{A}$. Hence π_x is a Fell point of $\hat{A}$, and A is a Fell algebra.

Conversely, we show that if X is not a Cartan G -space then A is not a Fell algebra. Some of the ideas come from the proof of [12, Proposition 4.2].

Suppose there is a point z in X with no wandering neighbourhood. To see that A is not a Fell algebra it is enough to produce an ideal I of the form $C_0(Y) \rtimes G$ which is not a Fell algebra. We may assume that A has bounded trace, because otherwise it is certainly not a Fell algebra.

To prove I is not a Fell algebra, we construct a net of irreducible representations $\{\pi_{x_W}\}$ of $\hat{I}$ parametrised by a directed family of neighbourhoods W of z , which converges to its unique cluster point π_z , and a positive element d of the Pedersen ideal of I such that $\mathrm{tr}(\pi_{x_W}(d))$ does not converge to $\mathrm{tr}(\pi_z(d))$. The minimality of the Pedersen ideal among dense ideals of I then implies that the ideal $\mathcal{S}(I)$ described in Equation $(\dagger)$ cannot be dense in I . It follows from [1, Corollary 3.7] that I is not a Fell algebra.

The construction of the element d and the net $\{\pi_{x_W}\}$ have to be carefully intertwined to ensure that we can compare the traces by finding appropriate eigenfunctions. We begin by building some functions in $C_c(G \times X)$ which will later be used to define d .

Let N be a compact neighbourhood of z in X and V_0 an open symmetric neighbourhood of e in G with compact closure V . Choose $f \in C_c(X)^+$ which is identically one on $V_0 \cdot N$.

For $x \in X$ we define $f_x(s) = f(s \cdot x)$. Let F be a symmetric compact subset of G such that $F \supseteq \text{supp}(f_z)$. Choose a non-negative function b in $C_c(G \times X)$ such that $b(r, x) = 1$ whenever $r \in VF^2V$ and $x \in \text{supp}(f)$. We can round off b so that $b(r, x) = 0$ whenever $(r, x) \notin V^2F^2V^2 \times M$, where M is a compact set containing the support of f . We would like to use the function $(t, x) \mapsto f(x)f(t^{-1} \cdot x)$ to define the element d , but this function will in general not be compactly supported; thus we will later multiply it by the function b .

Let $K = V^2F^2V^3F$. By [12, p. 61] there exists an open neighbourhood U of z , such that for every $x \in U$, the support of $f_x|_K$ is contained in $K \setminus V_0F$. To see this, let $\phi : G \times X \rightarrow X$ be the map $(s, x) \mapsto s \cdot x$. Observe that $(K \setminus V_0F) \times \{z\}$ is compact and is contained in the open subset $\phi^{-1}(X \setminus \text{supp}(f))$ of $G \times X$. For each $(s, z) \in (K \setminus V_0F) \times \{z\}$ there exist open neighbourhoods V_s of s and U_s of z such that $V_s \times U_s \subseteq \phi^{-1}(X \setminus \text{supp}(f))$. Choose a finite subcover $\{V_{s_i} \times U_{s_i}\}_{i=1}^n$ and set $U = \cap_{i=1}^n U_{s_i}$. Then

$$(K \setminus V_0F) \times \{z\} \subseteq \cup_{i=1}^n V_{s_i} \times U \subseteq \phi^{-1}(X \setminus M).$$

Thus for every $x \in U$, the function

$$f_x^1(s) := \begin{cases} f(s \cdot x) & \text{if } s \in K; \\ 0 & \text{if } s \notin V_0F \end{cases}$$

is well-defined and continuous on G . This function will be an eigenvector for the element d to be constructed. Without loss of generality we may suppose $U \subseteq N$.

Since no neighbourhood of z is wandering, for every neighbourhood W of z there are $x_W \in W$ and $s_W \notin K^{-1}K$ such that $s_W \cdot x_W \in W$. Thus we get a net $\{x_W\}$ directed by the neighbourhoods of z which converges to z , and by passing to a subnet if necessary, we may assume that this net is universal [9, 1.3.7] and that $x_W \in U$ for all W .

Because A has bounded trace and $\{\pi_{x_W}\}$ is a convergent net, the limit set of $\{\pi_{x_W}\}$ is discrete in the relative topology [6, Theorem 3]. The map $x \mapsto \pi_x := \text{Ind}_{\{e\}}^G(\epsilon_x)$ induces a homeomorphism of X/G onto $\hat{A}$, so there exists a neighbourhood V of $G \cdot z$ in X/G such that $G \cdot z$ is the unique limit point of the net $\{G \cdot x_W\}$ in V . We write $q : X \rightarrow X/G$ for the quotient map and let $Y = G \cdot (q^{-1}(V) \cap U)$. Since Y is open we may assume that $\{x_W\} \subseteq Y$. Observe that $\{\pi_{x_W}\}$ is the continuous image of a universal net, and hence is universal; thus $\{\pi_{x_W}\}$ is properly convergent.

We will show that the ideal $I = C_0(Y) \rtimes G$ of A is not a Fell algebra. Recall that our strategy to do this is to construct an element d in the Pedersen ideal of I such that $\{\text{tr}(\pi_{x_W}(d))\}$ does not converge to $\text{tr}(\pi_z(d))$. Let

$$B(t, x) = f(x)f(t^{-1} \cdot x)b(t^{-1}, x)\Delta(t^{-1})^{\frac{1}{2}} \quad \text{and} \quad D = \frac{1}{2}(B + B^*);$$

we will obtain the element d by pushing enough of $D^* * D$ into the Pedersen ideal of I .

The right Haar measure ν on G given by $\nu(E) = \mu(E^{-1})$ satisfies the hypothesis of [11, Lemma 4.14], so π_x is unitarily equivalent to the representation $M_x \rtimes V$ on $L^2(G, \nu)$ given by

$$V(t)\xi(s) = \Delta(t)^{\frac{1}{2}}\xi(t^{-1}s) \quad \text{and} \quad M_x(g)\xi(s) = g(s \cdot x)\xi(s)$$

for $\xi \in L^2(G, \nu)$ and $g \in C_0(X)$. Note that

$$\begin{aligned} M_x \rtimes V(B)\xi(s) &= \int_G B(t, s \cdot x) \Delta(t)^{\frac{1}{2}} \xi(t^{-1}s) d\mu(t) \\ &= \int_G f(s \cdot x) f(u \cdot x) b(us^{-1}, s \cdot x) \xi(u) d\nu(u), \end{aligned}$$

so that $M_x \rtimes V(B)$ is a Hilbert-Schmidt operator with kernel $(s, u) \mapsto f(s \cdot x) f(u \cdot x) b(us^{-1}, s \cdot x)$ in $C_c(G \times G)$.

By composing translation on the right by $(s_W)^{-1}$ with $f_{x_W}^1$ we see that

$$f_{x_W}^2(s) = \begin{cases} f(s \cdot x_W) & \text{if } s \in Ks_W; \\ 0 & \text{if } s \notin V_0Fs_W \end{cases}$$

is a well-defined continuous function on G whenever $x_W \in U \cap Y$. Note that $f_{x_W}^1$ and $f_{x_W}^2$ have disjoint supports because $s_W \notin K^{-1}K$. The $f_{x_W}^i$'s ($i = 1, 2$) are linearly independent eigenvectors for $M_{x_W} \rtimes V(D)$; as in the proof of [12, Lemma 4.4] one shows that

$$f(s \cdot x_W) f(u \cdot x_W) [b(us^{-1}, s \cdot x_W) + b(su^{-1}, u \cdot x_W)] f_{x_W}^i(u) = 2f_{x_W}^i(s) f(u \cdot x_W) f_{x_W}^i(u)$$

for all $s, u \in G$. The eigenvalue corresponding to $f_{x_W}^i$ is

$$\lambda_{x_W}^i = \int_G f(u \cdot x_W) f_{x_W}^i(u) d\nu(u) = \int_G f_{x_W}^i(u)^2 d\nu(u) = \int_G f_{x_W}^1(u)^2 d\nu(u) = \lambda_{x_W}^1.$$

If $x_W \in Y \cap U$ then $\lambda_{x_W}^1 \geq \nu(V_0) > 0$ because $u \in V_0$ implies $u \cdot x_W \in V_0 \cdot U \subseteq V_0 \cdot N$ and f was chosen to be identically one on $V_0 \cdot N$. It follows that if $x_W \in Y \cap U$ then $M_{x_W} \rtimes V(D^* * D)$ is a positive compact operator with eigenvalue $(\lambda_{x_W}^1)^2$ greater or equal to $a := \nu(V_0)^2$ of multiplicity at least 2.

On the other hand, $M_z \rtimes V(D)$ is the rank-one operator $f_z \otimes \bar{f}_z$ with eigenvalue $\lambda_z^1 = \|f_z\|^2 = \int_G f_z(s) d\nu(s) = \lim \lambda_{x_W}^1 > 0$. Thus $M_z \rtimes V(D^* * D)$ is a rank-one operator with eigenvalue $(\lambda_z^1)^2$.

Finally, we push enough of the operator $D^* * D$ into the Pedersen ideal of I using the description in [8, Theorem 5.5.1]. Let r be any non-negative function in $C_c((0, \infty))$ such that

$$r(t) = \begin{cases} 0 & \text{if } t < \frac{a}{3}; \\ 2t - \frac{2a}{3} & \text{if } \frac{a}{3} \leq t \leq \frac{2a}{3}; \\ t & \text{if } \frac{2a}{3} \leq t \leq \|D^* * D\|. \end{cases}$$

Now $d = r(D^* * D)$ is a positive element of the Pedersen ideal of I . If $x_W \in Y \cap U$ then $\text{tr}(M_{x_W} \rtimes V(d)) \geq 2(\lambda_{x_W}^1)^2 \rightarrow 2(\lambda_z^1)^2$ and $\text{tr}(M_z \rtimes V(d)) = (\lambda_z^1)^2$. The existence of such an element implies that I and hence A are not Fell algebras. $\square$

A C^* -algebra A always has a largest Fell ideal because the set of Fell points is open in $\hat{A}$.

Corollary 2. *Let (G, X) be a free second countable transformation group such that $C_0(X) \rtimes G$ is Type I. The largest Fell ideal of $C_0(X) \rtimes G$ corresponds to the open G -invariant subset $Y = \{x \in X : \exists \text{ a wandering neighbourhood of } x\}$ of X .*

Proof. If the action is free and $C_0(X) \rtimes G$ is Type I then every closed ideal is of the form $C_0(Y) \rtimes G$ for some open G -invariant subset of X by [3, Lemma 16]. Now apply Theorem 1 to $C_0(Y) \rtimes G$. $\square$

The following transformation group is interesting for several reasons. Its C^* -algebra is a Fell algebra which does not have continuous trace. In addition, it has many ideals with continuous trace but no largest continuous-trace ideal. Recall that a positive element a of a C^* -algebra A is a *continuous-trace element* if the function $\pi \mapsto \text{tr}(\pi(a))$ is finite and continuous on $\hat{A}$; the C^* -algebra has continuous trace if the linear span $\mathcal{M}(A)$ of these elements is a dense (hereditary) ideal of A . In particular, the closure of $\mathcal{M}(C_0(X) \rtimes G)$ in the example below is not a maximal continuous-trace ideal.

Example 3. This transformation group was described in [7, p. 298]. Let X be the strip $\{(x, y) : -1 \leq x \leq 1, y \in \mathbb{R}\}$ in the plane. The group action is by $G = \mathbb{R}$. Beyond the strip $-1 < x < 1$ the action moves a point according to

$$t \cdot (1, y) = (1, y + t) \quad \text{and} \quad t \cdot (-1, y) = (-1, y - t).$$

If $|x_0| < 1$ let $C_{(x_0, y_0)}$ be the vertical translate of the graph of the equation $y = \frac{x^2}{1-x^2}$ which passes through (x_0, y_0) . Define $t \cdot (x_0, y_0)$ to be the point (x, y) on $C_{(x_0, y_0)}$ such that the length of the arc of $C_{(x_0, y_0)}$ between (x_0, y_0) and (x, y) is $|t|$, and $x - x_0$ has the same sign as t . That is, (x_0, y_0) moves counter-clockwise along $C_{(x_0, y_0)}$ at unit speed.

A compact neighbourhood N in X is wandering if and only if it meets at most one boundary line $x = 1$ or $x = -1$; thus X is a Cartan G -space, but the action of G on X is not proper [7, p. 298]. Since the action is free, $C_0(X) \rtimes G$ does not have continuous trace, but it is a Fell algebra by Theorem 1.

The action of G on both $Y = \{(x, y) : -1 < x \leq 1, y \in \mathbb{R}\}$ and $Z = \{(x, y) : -1 \leq x < 1, y \in \mathbb{R}\}$ is free and proper, so that $I_1 = C_0(Y) \rtimes G$ and $I_2 = C_0(Z) \rtimes G$ are distinct maximal ideals which both have continuous trace.

We now identify the closure of the ideal of the continuous-trace elements. Let U be any compact wandering neighbourhood of $(1, 0)$ in X . Note that U wandering means that U does not meet the line $x = -1$. The saturation $G \cdot U$ is not closed because we can easily choose a sequence in $G \cdot U$ converging to $(-1, 0)$. The same is true for any compact wandering neighbourhood of $(1, y)$ or $(-1, y)$. By [3, Corollary 18] the closure of the continuous-trace ideal $\mathcal{M}(C_0(X) \rtimes G)$ is $C_0(W) \rtimes G$ where W consists of points (x, y) which have a wandering neighbourhood U such that $G \cdot U$ is closed in X . Thus $W = \{(x, y) : -1 < x < 1, y \in \mathbb{R}\}$ and $\overline{\mathcal{M}(C_0(X) \rtimes G)} = C_0(W) \rtimes G = I_1 \cap I_2$.

In [4] we characterise transformation groups whose C^* -algebras have bounded trace.

REFERENCES

1. Robert J. Archbold and Douglas W. B. Somerset. Transition probabilities and trace functions for C^* -algebras. *Math. Scand.*, 73:81–111, 1993.
2. Robert J. Archbold, Douglas W. B. Somerset, and John S. Spielberg. Upper multiplicity and bounded trace ideals in C^* -algebras. *J. Funct. Anal.*, 146:430–463, 1997.
3. Phillip Green. C^* -algebras of transformation groups with smooth orbit space. *Pacific J. Math.*, 140:71–97, 1977.
4. Astrid an Huef. *Transformation-group C^* -algebras with bounded trace*. PhD thesis, Dartmouth College, 1999.
5. Dale Husemoller. *Fibre Bundles*. Springer-Verlag, New York, second edition, 1975. Graduate Texts in Mathematics, No. 20.
6. Dragan Milićić. On C^* -algebras with bounded trace. *Glasnik Mat. Ser. III*, 8:7–22, 1973.

7. Richard S. Palais. On the existence of slices for actions of non-compact Lie groups. *Ann. of Math.*, 73:295–323, 1961.
8. Gert K. Pedersen. *C^* -algebras and their automorphism groups*. Academic Press, London, 1979.
9. Gert K. Pedersen. *Analysis now*. Springer-Verlag, New York, 1989.
10. Iain Raeburn. On crossed products and Takai duality. *Proc. Edinburgh Math. Soc.*, 31:321–330, 1988.
11. Dana P. Williams. The topology on the primitive ideal space of transformation group C^* -algebras and CCR transformation group C^* -algebras. *Trans. Amer. Math. Soc.*, 266:335–359, 1981.
12. Dana P. Williams. Transformation group C^* -algebras with continuous trace. *J. Funct. Anal.*, 41:40–76, 1981.

DEPARTMENT OF MATHEMATICS, DARTMOUTH COLLEGE, HANOVER, NH 03755, USA

E-mail address: `astrid.anhuef@dartmouth.edu`

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, UNIVERSITY OF DENVER, DENVER, CO 80208, USA

E-mail address: `astrid@cs.du.edu`